arXiv:hep-ph/0110358v1 27 Oct 2001

Confirmation of σ(450–600)-Meson in $\Upsilon' \to \Upsilon\pi\pi$ & Other $\pi\pi$-Production Processes

Muneyuki Ishida, Shin Ishida* Toshihiko Komada** and
Shin-Ichirou Matsumoto*

Department of Physics, Tokyo Institute of Technology
Tokyo 152-8551, Japan
**Atomic Energy Research Institute, College of Science and Technology*
Nihon University, Tokyo 101-0062, Japan
***Department of Engineering Science, Junior College Funabashi Campus, Nihon*
University, Funabashi 274-8501, Japan

(Received February 1, 2008)

Applying the effective amplitude, which is evidently consistent with general constraints from chiral symmetry, the $\pi\pi$ spectra in the relevant processes are analyzed, leading to a strong evidence for existence of the light σ meson. It is also pointed out that the $\pi\pi$ scattering process, which had been one of the main sources for PDG table for these many years, is, in principle, exceptionally difficult to investigate the property of σ-meson.

§1. Introduction

In the previous work referred to as I $^{1)}$, we have analyzed systematically the $\pi\pi$ production amplitudes in the various excited Υ and J/ψ decay processes, and reproduced successfully the experimental behaviors: The production amplitude $\mathcal{F}$ is given by a coherent sum of σ Breit-Wigner amplitude $\mathcal{F}_\sigma$ and direct 2π amplitude $\mathcal{F}_{2\pi}$ in the VMW method.

$$\mathcal{F} = \mathcal{F}_\sigma^I + \mathcal{F}_{2\pi}^I; \quad \mathcal{F}_\sigma^I = \frac{r_\sigma e^{i\theta_\sigma}}{m_\sigma^2 - s - i\sqrt{s}\Gamma_\sigma(s)}, \quad \mathcal{F}_{2\pi}^I = r_{2\pi} e^{i\theta_{2\pi}}, \qquad (1\cdot1)$$

where r_σ ($r_{2\pi}$) is production coupling constant of σ-state (2π-state) and $e^{i\theta_\sigma}$ ($e^{i\theta_{2\pi}}$) is a strong phase factor. Here $(r_\sigma, \theta_\sigma)$ and $(r_{2\pi}, \theta_{2\pi})$ are taken to be process-dependent, free parameters, since the $|\sigma\rangle$ state and $|2\pi\rangle$ state are, from quark physical picture $^{2)}$, independent bases of S-matrix with an independent vertex, in principle.

However, we must give special attention on the threshold behaviors. Because of the property of π meson as Nambu-Goldstone boson in the case of chiral symmetry breaking, the $|\mathcal{F}|^2$ was widely believed to be suppressed in the $\pi\pi$ threshold. In the $\pi\pi$ scattering, the observed spectrum is actually suppressed near the threshold. However, in the relevant excited-Υ decay processes this threshold suppression is also observed experimentally in $\Upsilon(2S \to 1S)$ and $\psi(2S \to 1S)$, while, in $\Upsilon(3S) \to \Upsilon(1S)\pi\pi$, the steep increase from the $\pi\pi$ threshold is observed.

In the following we examine the consistency of our results in I, especially the threshold behavior of $\Upsilon(3S \to 1S)$, with the constraint from chiral symmetry.

§2. Threshold Behavior of Production Amplitude and Chiral Symmetry

2.1. *Effective amplitude by linear σ model*

First we consider an effective chiral symmteric Lagrangian [3), 4)] including σ and π mesons, $\mathcal{L}^{(n)} = \xi^{(n)} \Upsilon'_\mu \Upsilon_\mu (\sigma^2 + \pi^2)$, where $\Upsilon'(\Upsilon)$ is the initial (final) $b\bar{b}$ quarkonium. Through the spontaneous breaking of chiral symmetry, σ acquires vacuum expectation value, $\langle\sigma\rangle_0 \equiv \sigma_0 = f_\pi$, and $\mathcal{L}^{(n)}$ is rewritten as $\mathcal{L}^{(n)} = \xi^{(n)} \Upsilon'_\mu \Upsilon_\mu (f_\pi^2 + 2 f_\pi \sigma' + \sigma'^2 + \pi^2)$, producing the coupling terms; $\mathcal{L}_\sigma = \xi_\sigma \Upsilon'_\mu \Upsilon_\mu \sigma'$ ($\xi_\sigma = 2 f_\pi \xi^{(n)}$), $\mathcal{L}_{\pi\pi} = \xi_{2\pi} \Upsilon'_\mu \Upsilon_\mu \pi^2$ ($\xi_{2\pi} = \xi^{(n)}$). The $\pi\pi$ production amplitude $\mathcal{F}^{L\sigma M}$ is given by sum of $\mathcal{F}_\sigma^{L\sigma M}$ and $\mathcal{F}_{2\pi}^{L\sigma M}$, canceling with each other in $O(p^0)$ level.

$$\mathcal{F}^{L\sigma M} = \mathcal{F}_\sigma^{L\sigma M} + \mathcal{F}_{2\pi}^{L\sigma M} = \frac{\xi_\sigma(-2 g_{\sigma\pi\pi})}{m_\sigma^2 - s} + 2\xi^{(n)} = 2\xi^{(n)} \frac{m_\pi^2 - s}{m_\sigma^2 - s}, \qquad (2\cdot1)$$

where the relation of SU(2)LσM, $g_{\sigma\pi\pi} = (m_\sigma^2 - m_\pi^2)/(2 f_\pi)$, is used, $s = -(p_1 + p_2)^2$ ($p_{1,2}$ being the pion momenta), and the factor of polarization vectors of initial and final $b\bar{b}$ quarkonia, $\epsilon(P') \cdot \tilde{\epsilon}(P)$, is omitted. The final amplitude takes $O(p^2)$ form, being consistent with the derivative coupling property of π meson. In the limit $p_{1\mu} \to 0_\mu$, $s \to m_\pi^2$ and $\mathcal{F}^{L\sigma M} \to 0$. Thus the amplitude has Adler 0, which leads to the threshold suppression in conformity with the observed threshold behavior in $\Upsilon(2S) \to \Upsilon(1S)\pi\pi$.

2.2. *Effective amplitude through intermediate glueball states*

Next we consider the effective amplitude through the intermediate production of the ground-state scalar and tensor glueballs. By using the framework of new classification scheme [5)], the effective interaction is given by $\mathcal{L}^G \approx (\xi^G / M'M) \partial_\mu \Upsilon'_\lambda \partial_\nu \Upsilon_\lambda (\partial_\mu \sigma \partial_\nu \sigma + \partial_\mu \pi \cdot \partial_\nu \pi)$, where $M'(M)$ is the mass of initial(final) Υ. ξ^G is coupling constant. It is notable that, after spontaneous symmetry breaking, it produces no σ-amplitude cancelling the direct 2π-amplitude. Then the production amplitude is given by

$$\mathcal{F}_{2\pi}^G = -\epsilon(P') \cdot \tilde{\epsilon}(P) (\xi^G / M'M)(P' \cdot p_1 P \cdot p_2 + P' \cdot p_2 P \cdot p_1)\ , \qquad (2\cdot2)$$

where $P'(P)$ is the momentum of initial(final) Υ.

The $\mathcal{F}_{2\pi}^G$ vanishes when $p_{1\mu} \to 0_\mu$. Thus, it has Adler 0, satisfying the general constraint from chiral symmetry. However, it does not vanish at $s = m_\pi^2$. In the relevant process, the Adler limit $p_{1\mu} \to 0_\mu$ corresponds to neglect $\Delta M \equiv M' - M$ in comparison with m_π. At $s = 4 m_\pi^2$ the pion four-momenta should be $p_{1\mu} = p_{2\mu}$, leading to $p_{10} = p_{20} \approx \Delta M/2 = 450\text{MeV} \gg m_\pi$ in $\Upsilon(3S) \to \Upsilon(1S)$ transition. Actually $\mathcal{F}_{2\pi}^G$ can be approximated as $\mathcal{F}_{2\pi}^G \approx -2\xi^G p_{10} p_{20}$, which is almost s-independent in all the physical region and has no zero close to the threshold. Correspondingly, there is no suppression near the threshold.

2.3. *Quantitative Analysis*

In the actual analysis we modify $\mathcal{F}^{L\sigma M}$ in Eq. (2·1), with inclusion of width of the σ-meson, $\Gamma_\sigma(s) (= g_\sigma^2 p_1(s)/(8\pi s))$ and of strong phase, as $\mathcal{F}_{\sigma+2\pi}^{\rm phen} = (\epsilon(P') \cdot \tilde{\epsilon}(P)) 2\xi^{(n)} e^{i\theta_\sigma} \frac{m_\pi^2 - s}{m_\sigma^2 - s - i\sqrt{s}\Gamma_\sigma(s)}$.



Fig. 1. Fit to the $\pi\pi$ invariant mass spectrum [4] by the amplitude explicitly consistent with chiral constraint: (a) $\Upsilon(3S) \to \Upsilon(1S)\pi\pi$, (b) $\Upsilon(3S) \to \Upsilon(2S)\pi\pi$ and (c) $\Upsilon(2S) \to \Upsilon(1S)\pi\pi$. The respective contributions from $\mathcal{F}^{\rm phen}_{\sigma+2\pi}$ and from $\mathcal{F}^{G}_{2\pi}$ are shown by dotted and dot-dashed lines.

The results of the fit by using the semi-phenomenological amplitude $\mathcal{F}^{\rm phen} \equiv \mathcal{F}^{\rm phen}_{\sigma+2\pi} + \mathcal{F}^{G}_{2\pi}$ are depicted in Fig. 1. The $\pi\pi$ mass spectra in $\Upsilon(2S) \to \Upsilon(1S)$, $\Upsilon(3S) \to \Upsilon(1S)$ and $\Upsilon(3S) \to \Upsilon(2S)$ transitions are fitted simultaneously with common values of m_σ and g_σ, leading to the σ-pole position, $\sqrt{s_{\rm pole}}(\approx m_\sigma - i\Gamma_\sigma/2) = (580\ {}^{+79}_{-30}) - i(190\ {}^{+107}_{-49})$, which is consistent with the ones obtained in our preceding works: [1] and the analysis [6] of $\pi\pi$ scattering phase shift. The total χ^2 is $\chi^2/(N_{\rm data} - N_{\rm param}) = 54.5/(74-11) = 0.87$. The $\mathcal{F}^{G}_{2\pi}$ contribution becomes dominant in $\Upsilon(3S) \to \Upsilon(1S)$ transition, and it explains the steep increase from the threshold.

Now we can see our treatment in the previous analysis [1] is to be consistent with general chiral constraints: Threshold suppression in $\Upsilon(2S) \to \Upsilon(1S)$, required from chiral symmetry, is reproduced phenomenologically by the cancellation between $\mathcal{F}^{I}_{\sigma}$ and $\mathcal{F}^{I}_{2\pi}$ of I (quoted in Eq. (1·1)). Dominant $\mathcal{F}^{G}_{2\pi}$-contribution, which is almost s-independent in all physical region, in $\Upsilon(3S) \to \Upsilon(1S)$ is reproduced mainly by constant $\mathcal{F}^{I}_{2\pi}$ of I .

2.4. *Features of $\pi\pi$ production amplitudes*

The $\pi\pi$ production processes have generally much the larger energy release ΔE than m_π. Thus, the momentum of emitted pion becomes large, and the derivative type amplitude, as Eq.(2·2), may play an important role. The Adler 0 generally does not imply the suppression at the small s region, and the $\pi\pi$ spectrum shows possibly



Fig. 2. Pole positions $\sqrt{s_{\rm pole}} \approx m_\sigma - i\Gamma_\sigma/2$ of σ and their error regions, obtained from the various $\pi\pi$-production processes: Υ decays [1] and [4], $p\bar{p} \to 3\pi^0$ [10], $D^- \to \pi^-\pi^+\pi^+$ [11] and $J/\psi \to \omega\pi\pi$ [9] are shown by the points with crosses. Respective regions (except for J/ψ decay) are surrounded by solid lines, dashed line and dot-dashed line.

steep increase from threshold. There is no mechanism cancelling the effect of σ production, and the direct σ-peak structure is expected to be observed. Actually,[*] in $J/\psi \to \omega\pi\pi$ decay [8), 9)], $p\bar{p} \to 3\pi^0$,[10] and $D^- \to \pi^-\pi^+\pi^+$ [11], where ΔE is very large, the σ-peak structure is observed directly. In the relevant Υ decay processes ΔE in $\Upsilon(3S \to 1S)$ is the largest, and only this process shows the steep increase from threshold.

On the contrary in the $\pi\pi$ scattering process, the pion momentum itself becomes small near threshold region, and the spectrum close to threshold is suppressed. Because of this chiral constraints, the σ-amplitude must be cancelled out by the non-resonant repulsive amplitude, and the direct σ-peak cannot be seen in $\pi\pi$-scattering, in principle.

§3. Concluding Remarks

Through the above discussion it proves not adequate to determine the pole-position of σ only through the $\pi\pi$-scattering. However, the present estimation of the σ pole position in PDG table has been done mainly through the analyses of $\pi\pi$ scattering. This is one of the reasons why the present label of σ, "f_0(400–1200) or σ", is largely uncertain and the σ is still regarded as controversial.

The σ-pole position should be determined regarding the $\pi\pi$-production processes with large energy release, which are free from the chiral constraints. The pole positions of σ obtained through various $\pi\pi$-production processes are shown in Fig. 2.

---

[*] Concerning the pp-central collision experiment $pp \to pp(\pi^0\pi^0)$ by GAMS [7], which also seems to suggest the σ-existence, see the comment in the talk by T. Komada, this conference [1].

They are almost process-independent, and in the range m_σ=450–600MeV. Basing on these results we conclude that *the existence of* σ(450–*600) is confirmed, and the present label of PDG table should be corrected accordingly.*

In this connection it is notable that, in this conference, a firm experimental evidence for existence of the $I = 1/2$ scalar κ-meson, to be the flavor partner of σ-meson, is reported. This κ is observed 11) as a peak in the $K\pi$-mass spectra in the decay process $D^- \to K^-\pi^+\pi^+$. This process is actually a $K\pi$-production process with the above mentioned large energy release, being free from the chiral constraints. The criticism 12) that no such pole is observed in $K\pi$-scattering process is disregarding the property of K meson as Nambu-Goldstone boson in the case of chiral symmetry breaking. This is the same situation as the direct σ-peak being not observed in $\pi\pi$-scattering. Actually by taking into account this constraint from chiral symmetry, the $K\pi$-scattering data were shown to be consistent with the exsitence of κ 13).

References

1) T. Komada, M. Ishida and S. Ishida, this conference; Phys. Lett. **B 508** (2001), 31.
2) S. Ishida, in proc. of Hadron97 in BNL, 1997, AIP conf. proc. 432; WHS99 in Frascati, 1999, Frascati Physics Series 15, 1999. M. Ishida, S. Ishida, T. Ishida, Prog. Theor. Phys. **99** (1998), 1031.
3) L. S. Brown and R. N. Cahn, Phys. Rev. Lett. **35** (1975), 1.
4) M. Ishida, S. Ishida, T. Komada and S. I. Matsumoto, Phys. Lett. **B 518** (2001), 47.
5) S. Ishida, this conference. S. Ishida, M. Ishida, T. Maeda, Prog. Theor. Phys. **104** (2000), 785.
6) S. Ishida et al., Prog. Theor. Phys. **95** (1996), 745; **98** (1997), 1005.
7) D. Alde et al., Phys. Lett. **B397** (1997), 350. T. Ishida, Doctor thesis, Univ. of Tokyo 1996.
8) J. E. Augustin et al., Nucl. Phys. **B320** (1989), 1.
9) K. Takamatsu, in proc. of Hadron97 at BNL, AIP conf. proc. 432, Upton NY 1997.
10) M. Ishida,T. Komada,S.Ishida,T. Ishida,K. Takamatsu,T. Tsuru, Prog. Theor. Phys. **104** (2000), 203.
11) C. Gobel, this conference.
12) S. N. Cherry and M. R. Pennington, Nucl. Phys. **A 688** (2001), 823.
13) S. Ishida, M. Ishida, T. Ishida, K. Takamatsu and T. Tsuru, Prog. Theor. Phys. **98** (1997), 621.